

Chapter 5

Nervous System Ageing

Claire Bénard and Maria Doitsidou

Abstract

In the face of ever-changing cellular environments during life and ageing, the nervous system ensures the coordination of behavior and physiology. Over time, however, the nervous system declines structurally and functionally, leading to age-related cognitive and behavioral decline in humans. Aspects of nervous system ageing are being studied using *C. elegans* as a model system. Here we review the age-related neuronal changes that occur at the structural, cellular and functional levels in normally ageing animals, as well as how these changes relate to lifespan in healthy ageing and in neurodegenerative conditions. Advances understanding the cellular mechanisms that result in neuronal decline in *C. elegans* will help identify cellular factors that protect the nervous system structure and function during normal ageing and in disease states. Ultimately, elucidating the molecular networks and cellular processes underlying the ageing of the nervous system will fuel research and design of interventions to improve human life at old age.

Keywords: *C. elegans*, ageing, aging, neuronal, neuron, nervous system, lifespan, longevity, behavior, decline, memory, learning, axon regeneration, neurodegeneration, insulin signaling, dietary restriction, mitochondria, proteostasis, protein aggregation

C. Bénard (✉)

Department of Neurobiology, University of Massachusetts Medical School, USA
e-mail: claire.benard@umassmed.edu

M. Doitsidou (✉)

Centre for Integrative Physiology, University of Edinburgh, UK.
e-mail: maria.doitsidou@ed.ac.uk

5.1 Introduction

Ageing precipitates alterations in the physiology of the nervous system, including age-related cognitive decline and an increased incidence of neurodegenerative diseases. Whereas age is known to be a strong determinant of these conditions, the etiology and molecular mechanisms leading to natural age-related neuronal deterioration are not well understood. Maintaining physiological functions with age depends on a continuous response to cellular stresses. General hallmarks of cellular ageing include DNA damage, loss of proteostasis, mitochondria dysfunction, autophagy impairment, loss of cytoskeletal integrity, nutrient sensing dysregulation, among others. The molecular pathways that regulate cellular aging are under intensive investigation and are reviewed elsewhere ([1, 2], chapters 6 to 12 of this volume). The nervous system is inevitably impacted by universal cellular processes that lead to cellular ageing, as well as by neuronal-specific factors.

C. elegans is a powerful system in which to elucidate the genetic networks and molecular pathways underlying neuronal aging. Its life cycle is fast, reaching adulthood in three days, generating its progeny in the following five days, and senescing in the following two weeks. Major conserved genetic determinants of lifespan have been elucidated [3, 4], enabling age manipulation in multiple ways. The worm's nervous system is simple, composed of exactly 302 identified neurons, which can be examined in exquisite detail in living animals at any point of their lives thanks to the worm's transparency and the ability to label specific neurons with fluorescent reporters. Further, the worm's entire neural circuitry has been defined, allowing one to probe neuronal structure and function in ageing animals. Importantly, the ease of genetic manipulation in the worm will fuel the identification of the genetic and molecular basis of neuronal aging. Given the extensive evolutionary conservation of the development and function of neurons between *C. elegans* and mammals, the worm offers the possibility to efficiently figure out fundamental principles by which the nervous system ages. Recent studies in *C.*

elegans have started to decipher the neuronal changes that accompany ageing and the factors that influence them, as we review below and summarize key findings in Figure 1 and Table 1.

5.2 Age-related structural and cellular changes in the nervous system

Similar to the healthy ageing human brain, the nervous system of *C. elegans* shows no neurodegeneration or gross deterioration during normal ageing [5-9]. Furthermore, the overall architecture of the nervous system is preserved throughout life ([5, 6, 8, 9], Bénard C, unpub.). However, like in humans, more subtle morphological neuronal changes do occur in ageing *C. elegans*. Hermaphrodites have been used in the studies reviewed here (except in the case of male mating behavior in section 5.4). Neuronal soma and axon diameter shrinks with age [5], and some neurons exhibit specific morphological changes, such as new branches along neuronal processes, axon swelling, axon waviness, defasciculation, new neurite-like extensions from the soma, and soma distortion ([6-9] and Bénard C, unpub.). These changes in neuronal morphology arise early during adulthood, progressively worsening in mid- (days 4 to 7) and old-aged animals ([6-9] and Bénard C, unpub.). The type of morphological change, age of onset, and frequency are highly neuron-type specific. Furthermore, the incidence and severity of these morphological changes vary among individual worms in isogenic populations that have been age-synchronized and co-cultured, suggesting that stochastic factors may influence these age-related neuronal changes.

Structural changes have been most extensively characterized in “gentle touch” mechanosensory neurons (ALM, PLM, AVM, PVM), each of which displays specific types of morphological changes. For instance, ectopic outgrowths appear from the soma of ALM by day 4 of adulthood, and new branches along the axon of PLM are frequent by day 8. Ectopic neurites sprouting from neuronal processes extend and retract dynamically [6, 7, 10]. Microtubule networks are disorganized in mechanosensory neurons with misshapen soma (ALM, [6]), and mitochondria are often located at the sites of ectopic neurites and swellings along the process [7]. The functional implications of these changes are unknown.

Other neurons also display age-related morphological changes, including branching from the soma of the dopaminergic neuron PDE from early adulthood onwards [7], defasciculation of cholinergic axons in the ventral nerve cord starting at day 6 of adulthood [6], axon beading of GABAergic neurons [6], and ectopic branches from GABAergic axons by day 5 [8]. Characterization of ageing in additional neuron types (e.g. other dopaminergic neurons, chemosensory neurons, interneurons, and motor neurons) extends the observation that age-related morphological changes are neuron-type specific, and widespread across the nervous system, but not ubiquitous ([9], Bernard C, unpub). It will be important to study a variety of neuronal types in mechanistic detail to forge a deeper understanding of the neuronal responses to age and elucidate the factors underlying the differential susceptibility of neurons to aging. Such analyses will provide insights into the basis of the selective neuronal vulnerability in neurodegenerative conditions in humans.

5.2.1 Synaptic deterioration in ageing neurons

As observed at the ultrastructural level, evidence of synaptic deterioration at day 15 of adulthood includes a decline in the synaptic vesicle numbers and a reduction in the size of presynaptic densities in the nerve cord and the nerve ring, which are sites of major synaptic contacts [7]. Synaptic vesicle density, observed using the fluorescently labeled synaptic vesicle protein RAB-3 GTPase, is also reduced in the presynaptic region of the motor neuron DA9 at day 18. Moreover, synaptic vesicle proteins (e.g. SNB-1/synaptobrevin and RAB-3 GTPase) ectopically accumulate in the dendritic and asynaptic axonal regions in ageing animals at day 12 and older (DA9 motor neurons, [11]). In addition, early endosomal membrane compartments (e.g. followed by RAB-5 GTPase, which are required for the formation and recycling of synaptic vesicles, are disorganized in ageing GABAergic motor neurons at day 10 [12]. Importantly, presynaptic release declines in motor neurons as early as day 7, and progressively worsens thereafter [13] (see also section 5.4). Age-related deterioration of synaptic organization, including an altered number of dendritic spines, has also been observed in mammals [14-16].

Axonal transport is key for synaptic maintenance during aging. At

the molecular level, genetic screening revealed two molecules that affect synaptic aging: the anterograde molecular motor UNC-104/KIF1A that transports synaptic vesicles and its regulator, the small GTPase ALR-1. Reduced function of UNC-104 accelerates synaptic deterioration and motor circuit dysfunction with age, whereas upregulation of UNC-104 improves synaptic function [11]. This highlights the importance of axonal transport in the maintenance of synaptic structural integrity throughout life.

5.2.2 Genetic factors that influence morphological ageing of the nervous system

Multiple conserved signaling pathways, including insulin signaling, dietary restriction, and mitochondria function, modulate the worm's lifespan (chapters 6 to 18, this volume). The insulin and insulin-like growth factor (IGF1) signaling pathway (IIS) is defined by *daf-2*, a homolog of the IGF-1 receptor (IGF1R) [17], which acts through the phosphatidylinositol 3-kinase PI3K kinase cascade. *daf-2* mutations increase lifespan [18] through changes in gene expression via activation of the downstream *daf-16* Forkhead box O (FOXO)-transcription factor, mutations in which shorten lifespan [19, 20]. *eat-2* encodes a subunit of a nicotinic acetylcholine receptor that functions in the pharynx [21]. Loss of function of *eat-2* serves as a genetic model of dietary restriction as it causes the worms to pump slower [21] and reduce their food intake, leading to a moderate increase in lifespan [22]. *clk-1* encodes the respiratory chain CoQ biosynthesis enzyme [23], and mutations in *clk-1* reduce respiration and extend lifespan [24].

As neurons undergo morphological changes with age, a fair expectation could be that long-lived mutants would delay neuromorphological aging, and conversely, short-lived mutants might accelerate neuronal changes. However, the relationship between lifespan pathways and age-related neuronal changes is complex, as only particular types of changes in certain neurons are affected by some but not all of the lifespan-altering mutations ([6-9], Bénard C, unpub.). For instance, studies show that whereas both *clk-1* and *eat-2* mutants have prolonged lifespans, neurite branching of mechanosensory neu-

rons is delayed in *clk-1* mutants, but not in *eat-2* mutants [8]. Also, *daf-2* mutants exhibit a delayed appearance of some of the branching defects [7, 8], but not of other age-related neuronal alterations ([7, 9], Bénard, unpub.). For example, ~8% of *daf-2* mutants exhibit novel defects at day 10 (e.g. branching from the ALM and PVM neurons), which is not seen in same-age or older wild-type animals ([7, 9] and Bénard, unpub.). Similarly, in the short-lived mutants *daf-16*, ALM soma outgrowth and PLMs with wavy axons are increased in early adulthood (day 2), but other aspects of neuronal morphology are unaffected and remain wild type ([7, 9] and Bénard, unpub.). Thus, lifespan genes differentially impact distinct types of neuronal changes, in a neuron-specific manner. Consistent with this notion, the organismal healthspan, as measured by locomotion, stress resistance, fat accumulation, muscle frailty, etc, does not always correlate with lifespan [25]. The separation of age-related morphological changes and lifespan is further revealed by tissue-specific manipulations of the *daf-2*/IGF1R pathway [8].

Other pathways that influence age-related morphological changes are the MAP kinase, heat shock stress response, and neuronal attachment pathways. The c-Jun terminal kinase JNK-1 and upstream kinases, JKK-1 and MEK-1, prevent the formation of ectopic neurite branching during ageing in a cell-autonomous manner [8]. In addition, the heat shock transcription factor HSF-1, which is under the control of the IIS pathway, is also required cell autonomously for maintaining neuronal integrity of ALM and PLM neurons. Finally, age-related defects in mechanosensory neurons are increased in the *mec-1* and *mec-5* mutants, in which the normal attachment of the touch neurons to the neighboring hypodermal cells is disrupted [6].

5.2.3 Maintenance of adult nervous system architecture

A number of genes of the immunoglobulin superfamily function to maintain neuronal architecture in *C. elegans* [26]. Some genes, such as the two-immunoglobulin domain containing proteins ZIG-4 and ZIG-3, act in early larval development to preserve the precise positioning of axons along the nerve cord. Other maintenance factors such as SAX-7, a homologue of L1CAM, and DIG-1, a large secreted protein required for basement membrane maintenance, play roles not only during larval development, but also during adulthood

where they maintain ganglia and nerve ring organization [27-32]. For instance, ganglia become disorganized in late larvae and adult *sax-7* mutants, in a way similar to the ganglia disorganization occurring in normally ageing wild-type adult animals, albeit earlier and more severely ([9], Bénard, unpub.) Furthermore, the two-immunoglobulin domain protein ZIG-10 is required continuously, including during adulthood, to maintain synapse density [33]. Such neuronal maintenance molecules, especially those mediating maintenance of the nervous system in adults, are likely to be neuro-protective during ageing ([9], Bénard, unpub.)

5.2.4 Subcellular changes in ageing neurons

As an organism ages, several features of senescence become apparent at the subcellular level, including alterations of organelle and cytoskeleton integrity, autophagic recycling, mitochondrial function and biogenesis, protein folding and homeostasis, telomere length, and transcriptional regulation, to name a few [2]. One of the challenges that neuronal cells face is to maintain adequate energy supply in distal neuronal processes, which they achieve by distributing mitochondria along axons and dendrites through specialized transport and anchoring [34]. Thus, processes that disturb the cytoskeletal network or mitochondrial function and transport can potentially affect healthy ageing and lead to neurodegenerative disease [35]. As mentioned above (section 5.2), such cellular events are affected in ageing *C. elegans* as microtubule networks become disorganized in neurons with age [6] and mitochondria localize at the base of age-related ectopic branches along neuronal processes [7].

The effect of ageing on *C. elegans* neuronal mitochondria in the cell body and processes of the mechanosensory neuron ALM was examined by Morsci *et al.* The frequency and distance of mitochondrial anterograde and retrograde transport progressively declines within the neuronal processes, starting already from the first day of adulthood, indicative of cytoskeletal transport decline [36]. Indeed, microtubules of mechanosensory neurons were shown to disorganize with age [6], and play a role in structural maintenance of neurons in the adult [37]. The size, density and stress resistance of mitochondria also change with age following a phasic pattern: first they increase during early adulthood (days 1 - 4), then they are maintained

at high levels in mid-adulthood (days 4 - 8), and finally they decline in later adulthood (days 8 - 15) [36]. The mitochondrial filamentous network becomes more complex and expansive in mid-adulthood whereas at later stages mitochondria exhibit ultrastructural abnormalities, e.g. loss of cristae structures [36]. Mitochondrial fragmentation was also observed in mechanosensory neurons and the ADF neurons [38]. By day 9 of adulthood, 50% of the ADF neurons exhibit fragmented mitochondria.

Mitochondrial changes are affected by lifespan mutations [36]. Mitochondrial fragmentation is attenuated in long-lived *daf-2*/IGF1R mutants, whereas it progresses more rapidly in short lived *hsf-1* mutants. *daf-2*/IGF1R mutants also have an elevated baseline oxidative stress level and do not exhibit decay in mitochondrial trafficking with age. Long-lived mutants *daf-2*, *eat-2* and overexpression of *sir-2.1* maintain a steady mitochondrial load during mid adulthood, in contrast to the elevated levels of same age wild-type animals [36]. Since compared to the wild type, long-lived mutants in general maintain a higher level of nervous system function at old age (see section 5.5), it appears that the mitochondrial profile of healthy neuronal ageing correlates with steady, rather than increased, mitochondrial content. How the interplay of mitochondrial biogenesis, degradation or fusion/fission dynamics brings about age-related mitochondrial changes and how these changes impact nervous system function, is under investigation in *C. elegans* and other models [39, 40].

5.2.5 Relevance to cellular changes in the human nervous system

In humans too, normal brain ageing is characterized by subtle changes in the morphology of specific neurons in selective brain regions [41, 42]. For instance, dendritic branching and length is enhanced in some hippocampal regions in aged individuals compared to young adults, and changes in dendritic spine and synapse number are observed in the ageing neocortex and hippocampus [41, 43, 44]. Despite the simplicity and short life of the *C. elegans* nervous system, its neurons -as described above- undergo age-related changes that parallel some neuronal changes in humans. Given the extensive evolutionary conservation of cellular processes between worms and humans, elucidating the mechanisms underlying the neuronal re-

sponses to ageing in *C. elegans* is expected to uncover conserved principles of neuronal aging.

5.3 Axon Regeneration and Ageing

Damaged axons have the ability to repair, which helps the nervous system to remain functional throughout life. In *C. elegans*, axons can be injured by laser axotomy and their regeneration examined with single-cell resolution. Severed axons frequently form a growth cone and regrow [45]. Multiple types of neurons, including mechanosensory neurons (ALM, PLM, AVM) and GABAergic motor neurons can regenerate, and the regenerative capacity differs among neuron types [46-49]. Similar to mammals, regrowth of injured axons in *C. elegans* is often misguided; nonetheless, regenerated axons appear to rewire -at least partly- into proper circuits, as demonstrated in worms that regain mobility after regeneration of their GABA motor neurons [46, 50].

Several molecular pathways that promote or inhibit axon regeneration have been discovered in *C. elegans* through genetic screening [51, 52]. Mechanisms of axon regeneration [53-57] include the PTEN and DLK-1 MAP kinase pathway and other MAP kinase pathways [52, 58-63], Notch signaling [55], microtubule regulators [51, 64, 65], and the IIS pathway [61]. Genetic analysis of axon regeneration has revealed that different neuron types share some regeneration genes, but have striking neuron-type-specific dependencies on other genes for axon regeneration.

5.3.1 Age-dependent decline of regeneration

Age is a strong determinant of a neuron's potential to drive axon repair. Young neurons regenerate damaged axons, but the regenerative ability of neurons quickly declines in early adulthood, worsening further with age [45]. Studies on the effect of age on axon regeneration have identified age-dependent mechanisms that regulate regenerative potential. In the mechanosensory neuron AVM, regeneration declines already during larval development and reaches stable levels that are sustained in adults. The pathway of miRNA *let-7* and its target gene *lin-41* regulates a switch from high capacity for

axon regrowth in early larvae when AVM develops, to low capacity for axon regrowth shortly after the developmental outgrowth of AVM is complete [66]. In contrast, the axon regrowth capacity of GABA motor neurons is high throughout larval stages and up to day 1 of adulthood, but steeply declines during adulthood (severely reduced by day five 5 and abolished by day 10) [58, 61]. This decline is a result of age-related deterioration in both axon initiation and axon elongation after injury. The insulin receptor DAF-2/IGF1R regulates this decline in GABA axon regeneration by inhibiting the *daf-16*/FOXO transcription factor and its downstream regulation of *dlk-1*/DLK and other genes of the DLK MAP kinase pathway [61]. Thus, *C. elegans* regulates the regenerative capacity of neurons in response to age.

The capacity of axons to regenerate in aging *C. elegans* does not directly correlate with lifespan, as not all long-lived mutants maintain regenerative capacity at old age. For instance, long-lived *eat-2* mutants and animals overexpressing *sir-2.1* have the same rates of regeneration as the wild type [61]. In contrast, loss of DAF-2/IGF1R function enhances regeneration of aged axons but not of young axons [61]. Neuron-specific expression of DAF-16/FOXO, which does not rescue lifespan, rescues axon regeneration in aged animals. Conversely, intestine-specific expression of DAF-16/FOXO, which rescues lifespan, does not rescue axon regeneration phenotypes in aged *daf-2* mutant animals. Thus, the role of the *daf-2/daf-16* pathway on axon regeneration is intrinsic to the nervous system and is uncoupled from its roles in lifespan regulation. The *C. elegans* adult neuronal IIS/FOXO transcriptome revealed the forkhead transcription factor FKH-9 as a IIS/FOXO target [67]. Loss of *fkh-9* impairs axon regeneration in aged *daf-2* mutants, and pan-neuronal expression of FKH-9 in *daf-2;fkh-9* mutants restored the regeneration phenotype, confirming its neuronal site of action [67].

5.3.2 Relevance to axon regeneration in ageing humans

During axon regeneration in *C. elegans* both age and neuron type determine a neuron's regenerative potential, partly because of specific dependencies on molecular pathways mediating axon regeneration. Similarly, age and neuron type strongly influence the regenerative capacity in humans. In adults, axons in the peripheral nervous

system regenerate, whereas axons in the central nervous system do not [68]. Intrinsic determinants of regeneration differ across the nervous system as well; for instance, removing PTEN greatly enhances optic and peripheral nerve regeneration, but has a modest effect on spinal cord axons [69-71]. These findings highlight the importance of studying diverse neuronal types in order to gain an understanding of regeneration, a goal that is achievable in the short term in *C. elegans* and that will inform research in mammals. Molecules identified in *C. elegans* to function in axon regeneration (e.g. PTEN and DLK), are conserved in mammals. Elucidating the mechanisms that regulate adult axon regeneration and the effect of age on neuronal regeneration will increase our understanding of how a neuron ages and inform approaches to treat injury and disease in humans.

5.4 Functional decline of the ageing nervous system

C. elegans is capable of versatile behaviors: it performs locomotion, rhythmic contractions for feeding known as pharyngeal pumping, defecation, egg-laying, and mating [72]. It also senses and responds to environmental cues including touch, odorants, temperature, and oxygen levels, and responds through the execution of behaviors such as the escape response, chemotaxis, and thermotaxis, to name a few [73]. As worms age, however, there is widespread behavioral decline [74]. The rate of locomotion slows down and eventually worms stop moving completely; in fact, a worm is considered dead when it fails to move in response to prodding. Measures of spontaneous locomotion (e.g. body bends, speed, turns, net displacement, trashing), as well as locomotion in response to a stimulus (e.g. chemotaxis and response to gentle touch), all decline with age [5, 74-81].

As a first step towards elucidating the causes for this behavioral decline, the ageing of both the muscles involved in the behavior, and of the neurons/neural circuits mediating the behavior needs to be examined. Body wall muscles, which power locomotion, have been found to progressively deteriorate with age in *C. elegans* starting at around day 10, and there is a clear correlation in individual animals between the severity of sarcopenia and the decline in locomotion [5]. This suggests that some of the age-related behavioral decline can be attributed to muscular deterioration. Several studies have tried to

tease apart muscle *vs.* neuronal contributions and although the primary cause has not yet been determined, there is clear evidence supporting a neuronal contribution to behavioral decline.

Clues about age-related decline in neuronal function came from pharmacological manipulations of the neuromuscular junction. Aged animals treated with the muscarinic agonist arecoline, which stimulates acetylcholine release from motor neurons, partially remedied age-related locomotion decline in day 8 and 10 animals. This raised the possibility that ageing affects neurotransmitter signaling at the neuromuscular junction [81]. Further pharmacological studies used the cholinergic agonist levamisole and the cholinesterase inhibitor aldicarb to stimulate body contractions throughout adulthood, including in very old worms that would otherwise be almost immobile (day 16); the findings indicated that presynaptic neuromuscular transmission declines after day 5 of adulthood [82]. Notably, 16-day old worms were capable of producing the same maximal contraction as young worms upon pharmacological stimulation of the neuromuscular junction, pointing to a neuronal synaptic contribution to the age-related locomotion decline [82]. Consistent with the notion that synaptic function deteriorates with age, synaptic terminal size and synaptic vesicle numbers decrease in old animals, and 18-day old animals that preserve higher synaptic integrity have better locomotion ability than same-aged animals with more deteriorated synapses [7].

Recently, electrophysiological studies using patch clamp recordings at the *C. elegans* neuromuscular junction provided direct functional evidence that the decline in presynaptic motor neuron function precedes muscle functional deficits [13]. The frequency of spontaneous neurotransmitter release from presynaptic motor neurons, measured as spontaneous post-synaptic currents, declines as early as day 5 of adulthood and progressively worsens with age, coinciding with locomotory decline. On the other hand, the amplitude of the postsynaptic currents recorded in muscles in response to presynaptic release did not change until day 11. Moreover, consistent with earlier pharmacological studies, the capacity of muscle contraction in response to levamisole did not decline before day 9. These experiments suggest that aging of the motor nervous system is an earlier

underlying reason for locomotory deterioration with age. Future studies will address the impact of the neuronal function decline on muscle deterioration, and the cellular and molecular basis leading to these functional changes in neurons.

C. elegans males perform mating behavior, which is a series of sensory and motor sub-behaviors to achieve copulation. The spicule intromission step of male copulation quickly deteriorates in adults and wild-type males' mating potency significantly decays by day 3 of adulthood (before age-related muscle deterioration), becoming impotent by day 5 [83]. Calcium imaging, pharmacological tests, and genetic manipulations showed that the decay in male mating results from increased excitability of the male sex muscles [84]. SIR-2.1, an ortholog of yeast SIR2 [85], is required to maintain mating potency, possibly by impacting metabolism at the level of glycolysis, fatty acid oxidation and oxidative stress responses, which in turn affect the excitability of the sex muscles [84].

Sensory perception-based behaviors also decline with age. While decay in locomotion with age certainly impairs these behaviors, age-related defects in the sensation and integration of stimuli also play a major role. Calcium imaging techniques, microfluidics and tools for neural circuit dissection have facilitated the study of neuronal function [86, 87]. Monitoring neuronal activity illuminates fine aspects of age-related changes in neuronal function that would be difficult to discern through behavioral analysis alone. For example, monitoring response to glycerol in the sensory neuron ASH revealed that calcium responses first increase in day 3-4 of adulthood, before they start decreasing in day 5 adults [88]. In a comprehensive analysis of odor-evoked neural signaling, Leinwand *et al.* identified a circuit of primary and secondary neurons that collectively encode benzaldehyde-evoked behavioral plasticity [89]. They find that the combinatorial circuit of odor sensing declines with age due to functional decline of the secondary but not the primary neurons, demonstrating that ageing differentially affects sensory neurons in the same circuit [89]. Whole-brain calcium imaging approaches in *C. elegans* [90, 91] will allow establishing temporal hierarchies of functional decline within neural circuits during ageing.

5.5. Learning and memory in aging

Learning and memory are fundamental biological processes that allow living organisms to respond and adapt to their environment. Memory decline in ageing is a well-documented phenomenon in humans [92] and a common feature across species [93]. Despite the simplicity of its nervous system, *C. elegans* exhibits behavioral plasticity and a range of well-characterized paradigms of short- and long-term memory [94]. These include examples of associative and non-associative memory, some of which have been studied in detail. Genetic pathways known to affect lifespan also affect learning and memory in different ways. In some cases, they play a role in the formation of memory itself; in other cases, they influence how fast memory declines during aging. Here, we focus on some of the well-characterized models of learning and memory in *C. elegans* to review the consequences of ageing on neuronal plasticity and the influence of lifespan-altering mutations on age-related decline.

5.5.1 Thermotaxis learning and memory in ageing

In a process known as thermotaxis, *C. elegans* has the ability to sense temperature and modify its behavior according to previous experience. It associates a past cultivation temperature with food and moves towards that temperature in search for food. Also, when found on a temperature gradient, it moves isothermally within the cultivation temperature. This behavior is called isothermal tracking and has characteristics of memory [95], for example it is CREB-dependent [96]. Thermotaxis learning declines with age, as isothermal tracking behavior becomes reduced by day 6 of adulthood and absent by day 14 among worms that retain mobility [97].

The IIS pathway, which regulates lifespan [18], affects associative learning behavior. In thermotaxis learning, aged animals of long-lived mutants *age-1/PI3K* and *daf-2/IGF1R* show enhanced isothermal tracking performance in young and old animals compared to the wild type. This enhancement consists of both a stronger association of temperature with food (as assessed by the number of animals performing isothermal tracking), as well as a delay in age-related decline of thermal learning [98]. This delay in age-related decline was not due to locomotion effects or simply a byproduct of increased lifespan: when Murakami *et al.* took into account the

physiological age (instead of the chronological), *age-1* mutants still had higher isothermal tracking behavior than the wild type [98]. In fact, there is a 210% extension in the period of high thermotaxis learning behavior in *age-1* mutants, compared to only 65% extension of the lifespan [98]. Moreover, expression of AGE-1/PI3K in AIY neurons in *age-1* mutants suppressed the *age-1* learning phenotype but not its longevity effects [98], thus dissociating lifespan from its effects on learning. The positive effect of *daf-2*/IGF1R and *age-1*/PI3K mutations on associative learning is *daf-16*/FOXO dependent. Interestingly, insulin peptide INS-1, the closest ortholog of human insulin, is required for the formation of the food-temperature association in a mechanism that acts antagonistically to the *daf-2*/IGF1R pathway [99].

In addition to the IIS pathway, thermotaxis learning is also affected by other longevity pathways. *eat-2* mutants, which extend lifespan through a dietary restriction mechanism, have enhanced thermotaxis learning in young adults but also delayed isothermal tracking decline with age [98]. Mitochondrial dysfunction also affects lifespan. Despite the complex roles of mitochondrial metabolism and ROS production in ageing [100], specific mutants of the electron transport chain are known to alter lifespan. Both *isp-1* (coding for the iron sulfur protein of mitochondrial respiratory complex III [101]) and *clk-1* (coding for a central enzyme in ubiquinone synthesis) mutants show increased thermotaxis learning behavior in young adult animals assessed by isothermal tracking [97]. The increased learning in *isp-1* mutants is *daf-16*/FOXO-dependent and it is abolished in *daf-16* mutants, despite the fact that the longevity effect of these mutants does not depend on *daf-16*/FOXO. Furthermore, *clk-1* mutants also delay age-related decline of isothermal tracking behavior. In contrast, short lived *gas-1* and *mev-1* mutants, defective for respiratory complex I and II, are more sensitive to oxidative stress [102, 103] and show decreased thermotaxis behavior, a phenotype rescued by treatment with antioxidants [97].

5.5.2 Positive olfactory associative learning and memory

C. elegans learns to associate volatile chemicals like butanone with food, and chemotaxes towards them [104]. When butanone is paired with food for a single training session (massed learning) it

produces a short-term associative memory (STAM). In contrast, when worms are subjected to multiple training sessions (spaced learning) they form a long-term associative memory (LTAM) that lasts between 16 and 24 hours [105]. LTAM formation declines with age, starting already at day 2 of adulthood and is abolished by day 5. LTAM deteriorates prior to the decline in olfactory learning, chemotaxis and motility [105], suggesting a higher sensitivity of LTAM in aging. Massed learning begins to decline already by day 3 and is completely lost by day 6 of adulthood. Spaced learning, which begins to decline on day 3, is lost by day 7.

Lifespan mutants affect LTAM and STAM in *C. elegans* both in young and older age. In young adults, *daf-2/IGF1R* mutants show three times longer STAM, although their learning rate is similar to wild type. Moreover, *daf-2* mutants show significantly longer LTAM, which remains active past 40 hours. Lastly, LTAM is established after fewer training sessions in *daf-2/IGF1R* mutants compared to wild-type animals. These memory improvements in young animals are *daf-16* dependent [105].

In older worms, *daf-2/IGFR* mutants retain their ability to learn for a longer time. At day 5 of adulthood, there is no significant loss in the formation of STAM. However, although learning is extended, LTAM is not improved in aged *daf-2/IGFR* mutants compared to the wild-type animals [105].

Dietary restriction affects positive olfactory associative memory differently than the IIS pathway. In young animals, *eat-2* mutants show no improvements in STAM compared to the wild type, whereas LTAM is reduced. In contrast, older *eat-2* animals show improved memory compared to wild type, and both STAM and LTAM persist for a longer period. Importantly, age-dependent memory loss can be alleviated if dietary restriction is imposed in adult worms [105].

5.5.3 Habituation (non-associative learning)

C. elegans also exhibits non-associative memory [94]. The best characterized examples are habituation to a mechanical stimulus and chemosensory habituation. These forms of adaptation can have

short- or long-term memory timescales depending on the training regime. Age-related changes in habituation have been reported: worms in their 6th and 8th day of adulthood habituate more rapidly to mechanical stimulus and show slower recovery from habituation than younger adults [106]. Timbers *et al.* tested adaptation to mechanical stimulus in middle-aged worms and showed that changes in habituation started at the peak of their reproductive age, as early as the 2nd day of adulthood [107]. This timeline is similar to the onset of changes in positive olfactory associative learning described above. In contrast to associative learning, the IIS pathway does not impact non-associative learning protocols involving chemosensory habituation [108].

5.5.4 Mechanisms of age-related learning and memory decline

Several forms of memory in *C. elegans*, for example olfactory STAM and LTAM, decline before any morphological neuronal changes become apparent [105, 109]. As a cautionary note, an analysis of age-related morphological changes of neurons that mediate learning and memory is still lacking. Thus, it seems that changes at the molecular level, which precede obvious morphological defects, are responsible for memory decline [105, 109]. Indeed, LTAM deterioration with age in both the wild type and longevity mutants correlates tightly with *crh-1*/CREB expression levels [105]. This correlation appears to be conserved in mammals: the levels of CREB in the brain are predictive of spatial memory decline in aged rats [110] and overexpression of CREB in the hippocampus attenuates spatial memory impairment during ageing [111].

Parallels can be drawn between memory and synapse decline during ageing in the wild type and lifespan mutants. The complex synaptic machinery required for memory formation is well documented [112]. *C. elegans* research has revealed a correlation between synapse deterioration and STAM decline with age [11]. At the molecular level, the anterograde kinesin motor UNC-104/KIF1A, which transports synaptic vesicles along axons, is required for STAM maintenance in aging. UNC-104/KIF1A levels are reduced with age in the wild type but maintained in *daf-2* mutants in a *daf-16*/FOXO-dependent manner [11].

The examples above demonstrate that a reduction in the IIS pathway promotes positive associative learning and memory in ageing *C. elegans*, through activation of the DAF-16/FOXO transcription factor. Neuron-specific transcriptome analysis of DAF-16/FOXO targets in *daf-2/IGF1R* mutants revealed a landscape of DAF-16/FOXO-dependent regulators of short-term memory extension, distinct from previously identified targets in other tissues. This analysis showed that some of the DAF-16/FOXO neuronal targets that extend memory in *daf-2/IGF1R* mutants also regulate memory in the wild type [67]. Thus, IIS pathway-dependent memory extension is due to augmentation or maintenance of the molecular machinery that regulates memory in the wild type, rather than the activation of an alternative mechanism. Similarly in mouse, FOXO6 is highly expressed in adult hippocampus and is required for memory consolidation by regulating the expression of genes responsible for synaptic function [113].

Among the DAF-16/FOXO targets that are upregulated in *daf-2/IGF1R* mutants at the whole worm level is FKH-9. It was shown that FKH-9 is required in the neurons for memory enhancement in *daf-2* mutants and in the somatic cells for lifespan extension [113]. Molecular characterization of the tissue-specific transcriptional programs that regulate longevity or neuronal function, combined with an analysis of the conservation of these programs across phylogeny, will facilitate a more complete understanding of nervous system ageing.

5.5.6 Relevance to learning and memory in humans

As described above, learning and memory decline in *C. elegans* becomes apparent in ageing animals in early adulthood. Recent studies indicate that cognitive decline in humans might start as early as the fourth decade of life [114]. *C. elegans* findings demonstrated the positive role of IIS pathway reduction and dietary restriction in delaying the decline of learning and memory. In humans, IIS plays physiological roles in various regions of the central nervous system, regulating neuronal function, including learning and memory. However, impaired insulin signaling and resistance observed in age-related diseases complicates the role of IIS in the human nervous

system [115]. Clarifying discrepancies will delineate its exact mechanisms of action in the ageing human brain.

Dietary restriction in humans has beneficial effects on brain structure and function [116]. Brain regions most vulnerable to the ageing process, namely frontal and medial temporal lobes, are negatively affected in obese individuals [117]. Excessive energy intake and elevated levels of blood glucose and fatty acids negatively impacts cognition [118]. In contrast, caloric restriction in elderly adults was shown to improve memory [119]. Recent evidence suggests that memory enhancement is a result of a negative energy balance (weight loss phase) rather than an effect of low weight maintenance [116]. Consequently, intermittent fasting or other interventions that mimic the effects of dietary restriction [120, 121] could prove more beneficial to healthy cognitive ageing than constant dietary restriction. *C. elegans* research has indeed demonstrated that intermittent fasting extends lifespan through the action of a small GTPase, RHEB-1, via the IIS pathway. Inhibition of RHEB-1 successfully mimicked the effects of caloric restriction. Thus, this and similar studies in *C. elegans* [122] can open the way to develop mimetics of dietary restriction to improve lifespan, health and cognitive function without the potential negative effects of caloric restriction.

5.6 Neurodegenerative diseases

Age is the leading risk factor for neurodegenerative disease [42, 123, 124]. This suggests that cellular changes occurring during ageing increase the vulnerability of neurons to such conditions. Ageing cells show increased levels of oxidative, metabolic and ionic stress that result in the accumulation of dysfunctional organelles, damaged proteins and DNA. Failure of neurons to adapt to such stresses leads to neuronal dysfunction and susceptibility to neuronal degeneration. Understanding the molecular mechanisms underlying age-related neurodegenerative diseases and identifying neuroprotective strategies is a major focus of modern medical research.

5.6.1 *C. elegans* models of neurodegenerative disease

Besides the general advantages of *C. elegans* as a model organism (discussed in section 5.1 and chapter 1 of this volume) additional characteristics make it particularly suitable for studying human neu-

rodenerative disease: *C. elegans* tolerates nervous system defects very well, as most of its neurons are dispensable for survival and reproduction in laboratory conditions [125, 126]. Furthermore, the majority of human genes implicated in monogenic forms of neurodegenerative disease have conserved *C. elegans* orthologs. When an ortholog does not exist, then expression of the human gene has often been used to reproduce disease phenotypes. Lastly, the power of genetic screens in *C. elegans* renders it ideal for rapid, genome-wide discovery of disease modifiers. Therefore, *C. elegans* has been extensively used to model and study neurodegenerative diseases [127-129].

Here we briefly summarize some examples of human neurodegenerative diseases studied in *C. elegans*, several of which are characterized by protein misfolding and aggregation (see also chapter 12). Parkinson's disease (PD) is an age-related movement disorder, accompanied by loss of dopamine neurons. *C. elegans* models of PD include overexpression of human α -synuclein in muscles or neurons, mutations in orthologs of human Parkinsonism (PARK) genes, and environmental toxins (such as paraquat, rotenone, MPTP, MPP+ and 6-OHDA) [130, 131]. Alzheimer's disease (AD) results in a progressive loss of cognitive function and is characterized by extracellular deposits of amyloid β (A β) and intracellular aggregates of microtubule associated protein *tau* (*MAPT*). *C. elegans* models of AD include overexpression of human A β peptides or tau [132-134]. Huntington's disease (HD) is a fatal neurodegenerative disorder caused by polyglutamine (polyQ) repeat expansion in huntingtin (*HTT*), and the corresponding *C. elegans* models express fragments of the human protein with various lengths of polyQ repeats [135]. Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease affecting motor neurons, which is caused by diverse genetic mutations. Most models of ALS in *C. elegans* relay on overexpression of wild-type and mutated forms of the causative genes *superoxide dismutase 1* (*SOD1*), *TAR DNA binding protein 43* (*TDP-43*) and *Fused-in-sarcoma* (*FUS*) [136]. Finally, a number of *C. elegans* models exist for studying neuronal channelopathies and excitotoxic cell death [137-139].

Modifier screens on the above models have contributed important insights into the understanding of neurodegenerative disease and revealed numerous general and disease-specific modifiers conserved in other organisms. Studies on *C. elegans* models of PD led to the discovery of several neuroprotective mechanisms; for instance, overexpression of the chaperone *TOR2/TorsinA*, the lysosomal P-ATPase *catp-6/ATP13A2*, or human Cathepsin D were shown to ameliorate aspects of α -synuclein toxicity [140-142] and the glycolytic enzyme GPI-1/GPI was identified as a conserved modifier of dopaminergic degeneration [143]. Genetic screens in *C. elegans* models of AD led to the discovery of many disease modifiers, e.g. orthologs of human kinases such as *kin-18/TAOK1* and *sgg-1/GSK3 β* , and chaperone stress response molecules (such as *xbp-1/XBP1*, *hsp-2/HSPA2*, *hsf-1/HSF1* and *chn-1/CHN1*) were identified as key regulators of *tau* toxicity [144]. A long list of similar discoveries has confirmed the validity of *C. elegans* as a model to study neurodegenerative disease. Key discoveries stem not only from research on models of disease but also from studies that enhanced our understanding of the normal function of disease-associated genes in healthy situations. A comprehensive review of these findings is beyond the scope of this chapter.

5.6.2 Ageing pathways and neurodegenerative disease

Although familial cases of neurodegenerative diseases in humans start earlier in life, the vast majority of cases for numerous neurodegenerative diseases are sporadic and manifest during the seventh decade or later, making ageing the major risk factor [145]. Is it then possible to prevent or delay such conditions by delaying ageing? Several studies in *C. elegans* show that manipulating longevity pathways affects pathological manifestations in models of neurodegenerative disease [146].

In *C. elegans* models of neurodegenerative disease, the IIS pathway modulates protein aggregation and toxicity. Reducing insulin signaling in *C. elegans* models of HD and AD alleviated polyglutamine [147, 148] and A β toxicity [149, 150], respectively. Similar protective effects of IIS reduction were reported in an ALS model of SOD-1 aggregation [151] and in an α -synuclein overexpression

model [143]. These results of reduced proteotoxicity across disease models are consistent with findings that overall protein insolubility and aggregation (an inherent part of normal ageing in *C. elegans* and other animals) is also alleviated by reduction in IIS signaling [152]. The mechanism of the *daf-2* mediated protection depends on the action of DAF-16/FOXO and HSF-1. These transcription factors have in fact opposing protective effects in AD models, with HSF-1 promoting disaggregation of A β , and DAF-16 promoting the active aggregation of toxic oligomers to less toxic forms [149]. Longevity manipulations through dietary restriction also suppress both polyglutamine and A β toxicity in *C. elegans* [153] and protect dopaminergic neurons from degeneration in a 6-OHDA model [154]. Finally, consistent with major longevity pathways modulating neurodegenerative disease, germ-cell ablation also attenuated polyglutamine toxicity in a DAF-16/FOXO and HSF-1 dependent manner [155].

The protective link between reduced IIS signaling and neuronal proteotoxicity was shown to be conserved in mice [156, 157], validating the relevance of the *C. elegans* findings. It also extends to neurological conditions that are not based on proteotoxic aggregation, for example hypoxia-induced ischemic stroke [158]. Similarly, the protective role of dietary restriction is conserved in mouse models of β -amyloid neuropathy [159] and in non-human primate model of PD [160]. Importantly, the effects exerted by ageing manipulations can be, at least in some cases, uncoupled from the extension of lifespan. This was demonstrated in mice by reducing IIS later in life when it can no longer extend the lifespan, a manipulation that nevertheless protected from A β toxicity [161]. The interplay of longevity pathways and disease has obvious implications in medical and lifestyle interventions, which could potentially delay or ameliorate devastating age-related disorders.

5.7 Concluding remarks

Studies in *C. elegans* have contributed to our understanding of the molecular machineries that protect the nervous system structure and function during normal ageing, following injury, and in disease states. One striking property of the ageing nervous system in *C. elegans*, which is also common in humans, is the differential suscepti-

bility of neurons to age. Moreover, whereas the decline of overall nervous system function is delayed in lifespan-extending mutations, specific aspects of neuronal deterioration are not, and specific neuron types are impacted differently by these lifespan mutations. The precise descriptions of age-related nervous system changes reviewed here constitute the basis for future mechanistic studies of neuronal ageing. Longitudinal analysis and the development of more tools to measure diverse aspects of neuronal ageing simultaneously will help establish relationships between age-related changes and identify the genetic and environmental factors underlying ageing of the nervous system. Ultimately, what we learn about the mechanisms influencing neuronal ageing will facilitate the development of therapeutic interventions to help improve the human condition in old age.

5.8 Acknowledgements

We thank Arantza Barrios, Emanuel K. Busch and Cassandra Blanchette for feedback on the manuscript. Research in the lab of Dr. Maria Doitsidou is supported by the Norwegian Research Council and the Wellcome Trust, UK. Research in the lab of Dr. Claire Bénard is supported by grant R01 AG041870-01A1 from the National Institutes of Health of the USA to C.B., the Ellison Medical Foundation New Scholar Aging Award to C.B., and the American Federation for Aging Research Award to C.B..

Dedicated to the memory of Muhammad Ali (January 17, 1942 - June 3, 2016).

References

1. López-Otín C, Blasco MA, Partridge L, et al (2013) The hallmarks of aging. *Cell* 153:1194–1217. doi: 10.1016/j.cell.2013.05.039
2. DiLoreto R, Murphy CT (2015) The cell biology of aging. *Mol Biol Cell* 26:4524–4531. doi: 10.1091/mbc.E14-06-1084
3. Kenyon CJ (2010) The genetics of ageing. *Nature* 464:504–512. doi: 10.1038/nature08980
4. Hekimi S, Guarente L (2003) Genetics and the specificity of the aging process. *Science* 299:1351–1354. doi: 10.1126/science.1082358

5. Herndon LA, Schmeissner PJ, Dudaronek JM, et al (2002) Stochastic and genetic factors influence tissue-specific decline in ageing *C. elegans*. *Nature* 419:808–814. doi: 10.1038/nature01135
6. Pan C-L, Peng C-Y, Chen C-H, McIntire S (2011) Genetic analysis of age-dependent defects of the *Caenorhabditis elegans* touch receptor neurons. *Proc Natl Acad Sci USA* 108:9274–9279. doi: 10.1073/pnas.1011711108
7. Toth ML, Melentijevic I, Shah L, et al (2012) Neurite sprouting and synapse deterioration in the aging *Caenorhabditis elegans* nervous system. *J Neurosci* 32:8778–8790. doi: 10.1523/JNEUROSCI.1494-11.2012
8. Tank EMH, Rodgers KE, Kenyon C (2011) Spontaneous age-related neurite branching in *Caenorhabditis elegans*. *J Neurosci* 31:9279–9288. doi: 10.1523/JNEUROSCI.6606-10.2011
9. Khandekar A (2015) Age-related Changes in the Neuronal Architecture of *Caenorhabditis elegans*. Doctoral thesis, Bénard Laboratory, University of Massachusetts Medical School, 2015.
10. Peng C-Y, Chen C-H, Hsu J-M, Pan C-L (2011) *C. elegans* model of neuronal aging. 4:696–698.
11. Li L-B, Lei H, Arey RN, et al (2016) The Neuronal Kinesin UNC-104/KIF1A Is a Key Regulator of Synaptic Aging and Insulin Signaling-Regulated Memory. *Curr Biol* 26:605–615. doi: 10.1016/j.cub.2015.12.068
12. Sann SB, Crane MM, Lu H, Jin Y (2012) Rabx-5 regulates RAB-5 early endosomal compartments and synaptic vesicles in *C. elegans*. *PLoS ONE* 7:e37930. doi: 10.1371/journal.pone.0037930
13. Liu J, Zhang B, Lei H, et al (2013) Functional aging in the nervous system contributes to age-dependent motor activity decline in *C. elegans*. *Cell Metab* 18:392–402. doi: 10.1016/j.cmet.2013.08.007
14. Valdez G, Tapia JC, Kang H, et al (2010) Attenuation of age-related changes in mouse neuromuscular synapses by caloric restriction and exercise. *Proc Natl Acad Sci USA* 107:14863–14868. doi: 10.1073/pnas.1002220107
15. Chen S, Hillman DE (1999) Dying-back of Purkinje cell dendrites

- with synapse loss in aging rats. *J Neurocytol* 28:187–196. doi: 10.1023/A:1007015721754
16. Rogers J, Zornetzer SF, Bloom FE, Mervis RE (1984) Senescent microstructural changes in rat cerebellum. *Brain Research* 292:23–32. doi: 10.1016/0006-8993(84)90886-2
 17. Kimura KD, Tissenbaum HA, Liu Y, Ruvkun G (1997) *daf-2*, an Insulin Receptor-Like Gene That Regulates Longevity and Diapause in *Caenorhabditis elegans*. *Science* 277:942–946. doi: 10.1126/science.277.5328.942
 18. Kenyon C, Chang J, Gensch E, et al (1993) A *C. elegans* mutant that lives twice as long as wild type. *Nature*
 19. Lin K, Dorman JB, Rodan A, Kenyon C (1997) *daf-16*: An HNF-3/forkhead Family Member That Can Function to Double the Life-Span of *Caenorhabditis elegans*. *Science* 278:1319–1322. doi: 10.1126/science.278.5341.1319
 20. Ogg S, Paradis S, Gottlieb S, et al (1997) The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in *C. elegans*. *Nature* 389:994–999. doi: 10.1038/40194
 21. McKay JP (2004) *eat-2* and *eat-18* Are Required for Nicotinic Neurotransmission in the *Caenorhabditis elegans* Pharynx. *Genetics* 166:161–169. doi: 10.1534/genetics.166.1.161
 22. Lakowski B, Hekimi S (1998) The genetics of caloric restriction in *Caenorhabditis elegans*. *Proceedings of the National ...*
 23. Ewbank JJ, Barnes TM, Lakowski B, et al (1997) Structural and Functional Conservation of the *Caenorhabditis elegans* Timing Gene *clk-1*. *Science* 275:980–983. doi: 10.1126/science.275.5302.980
 24. Lakowski B, Hekimi S (1996) Determination of Life-Span in *Caenorhabditis elegans* by Four Clock Genes. *Science* 272:1010–1013. doi: 10.1126/science.272.5264.1010
 25. Bansal A, Zhu LJ, Yen K, Tissenbaum HA (2015) Uncoupling lifespan and healthspan in *Caenorhabditis elegans* longevity mutants. *Proc Natl Acad Sci USA* 112:E277–86. doi: 10.1073/pnas.1412192112

26. Bénard C, Hobert O (2009) Chapter 6 Looking Beyond Development: Maintaining Nervous System Architecture. In: *Development of Neural Circuitry*. Elsevier, pp 175–194
27. Bénard CY, Boyanov A, Hall DH, Hobert O (2006) DIG-1, a novel giant protein, non-autonomously mediates maintenance of nervous system architecture. *Development* 133:3329–3340. doi: 10.1242/dev.02507
28. Burket CT, Higgins CE, Hull LC, et al (2006) The *C. elegans* gene dig-1 encodes a giant member of the immunoglobulin superfamily that promotes fasciculation of neuronal processes. *Dev Biol* 299:193–205.
29. Johnson RP, Kramer JM (2012) Neural Maintenance Roles for the Matrix Receptor Dystroglycan and the Nuclear Anchorage Complex in *Caenorhabditis elegans*. *Genetics* 190:1365–1377. doi: 10.1534/genetics.111.136184
30. Sasakura H, Inada H, Kuhara A, et al (2005) Maintenance of neuronal positions in organized ganglia by SAX-7, a *Caenorhabditis elegans* homologue of L1. *EMBO J* 24:1477–1488. doi: 10.1038/sj.emboj.7600621
31. Pocock R, Bénard CY, Shapiro L, Hobert O (2008) Functional dissection of the *C. elegans* cell adhesion molecule SAX-7, a homologue of human L1. *Molecular and Cellular Neuroscience* 37:56–68.
32. Wang X, Kweon J, Larson S, Chen L (2005) A role for the *C. elegans* L1CAM homologue lad-1/sax-7 in maintaining tissue attachment. *Dev Biol* 284:273–291. doi: 10.1016/j.ydbio.2005.05.020
33. Cherra SJ III, Jin Y (2016) A Two-Immunoglobulin-Domain Transmembrane Protein Mediates an Epidermal-Neuronal Interaction to Maintain Synapse Density. *Neuron* 89:325–336.
34. Sheng Z-H (2014) Mitochondrial trafficking and anchoring in neurons: New insight and implications. *J Cell Biol* 204:1087–1098. doi: 10.1083/jcb.201312123
35. Lionaki E, Markaki M, Palikaras K, Tavernarakis N (2015) Mitochondria, autophagy and age-associated neurodegenerative diseases: New insights into a complex interplay. *Biochim Biophys Acta* 1847:1412–1423. doi: 10.1016/j.bbabbio.2015.04.010

36. Morsci NS, Hall DH, Driscoll M, Sheng Z-H (2016) Age-Related Phasic Patterns of Mitochondrial Maintenance in Adult *Caenorhabditis elegans* Neurons. *J Neurosci* 36:1373–1385. doi: 10.1523/JNEUROSCI.2799-15.2016
37. Chew YL, Fan X, Götz J, Nicholas HR (2013) PTL-1 regulates neuronal integrity and lifespan in *C. elegans*. *Journal of Cell Science* 126:2079–2091. doi: 10.1242/jcs.jcs124404
38. Jiang H-C, Hsu J-M, Yen C-P, et al (2015) Neural activity and CaMKII protect mitochondria from fragmentation in aging *Caenorhabditis elegans* neurons. *Proc Natl Acad Sci USA* 112:8768–8773. doi: 10.1073/pnas.1501831112
39. Palikaras K, Lionaki E, Tavernarakis N (2015) Coordination of mitophagy and mitochondrial biogenesis during ageing in *C. elegans*. *Nature* 521:525–528. doi: 10.1038/nature14300
40. Lionaki E, Markaki M, Palikaras K, Tavernarakis N (2015) Mitochondria, autophagy and age-associated neurodegenerative diseases: New insights into a complex interplay. *Biochim Biophys Acta* 1847:1412–1423. doi: 10.1016/j.bbabi.2015.04.010
41. Burke SN, Barnes CA (2006) Neural plasticity in the ageing brain. *Nat Rev Neurosci* 7:30–40. doi: 10.1038/nrn1809
42. Yankner BA, Lu T, Loerch P (2008) The aging brain. *Annu Rev Pathol* 3:41–66. doi: 10.1146/annurev.pathmechdis.2.010506.092044
43. Esther A Nimchinsky, Bernardo L Sabatini A, Svoboda K (2003) Structure and Function of Dendritic Spines. <http://dxdoiorg/101146/annurevphysiol64081501160008> 64:313–353. doi: 10.1146/annurev.physiol.64.081501.160008
44. Hof PR, Morrison JH (2004) The aging brain: morphomolecular senescence of cortical circuits. *Trends Neurosci* 27:607–613.
45. Hammarlund M, Jin Y (2014) Axon regeneration in *C. elegans*. *Curr Opin Neurobiol* 27:199–207.
46. Yanik MF, Cinar H, Cinar HN, et al (2004) Neurosurgery: functional regeneration after laser axotomy. *Nature*
47. Yanik MF, Cinar H, Cinar HN, Gibby A (2006) Nerve regeneration in *Caenorhabditis elegans* after femtosecond laser axotomy.

48. Wu Z, Ghosh-Roy A, Yanik MF, et al (2007) *Caenorhabditis elegans* neuronal regeneration is influenced by life stage, ephrin signaling, and synaptic branching. *Proc Natl Acad Sci USA* 104:15132–15137. doi: 10.1073/pnas.0707001104
49. Gabel CV, Antoine F, Chuang C-F, et al (2008) Distinct cellular and molecular mechanisms mediate initial axon development and adult-stage axon regeneration in *C. elegans*. *Development* 135:1129–1136. doi: 10.1242/dev.013995
50. Bejjani El R, Hammarlund M (2012) Notch Signaling Inhibits Axon Regeneration. *Neuron* 73:268–278. doi: 10.1016/j.neuron.2011.11.017
51. Chen L, Wang Z, Ghosh-Roy A, et al (2011) Axon Regeneration Pathways Identified by Systematic Genetic Screening in *C. elegans*. *Neuron* 71:1043–1057.
52. Nix P, Hammarlund M, Hauth L, et al (2014) Axon regeneration genes identified by RNAi screening in *C. elegans*. *J Neurosci* 34:629–645. doi: 10.1523/JNEUROSCI.3859-13.2014
53. Hilliard MA (2009) Axonal degeneration and regeneration: a mechanistic tug-of-war. *J Neurochem* 108:23–32. doi: 10.1111/j.1471-4159.2008.05754.x
54. Wang Z, Jin Y (2011) Genetic dissection of axon regeneration. *Curr Opin Neurobiol* 21:189–196.
55. Rachid El Bejjani MH (2012) Neural Regeneration in *Caenorhabditis elegans*. *Annu Rev Genet* 46:499–513. doi: 10.1146/annurev-genet-110711-155550
56. Chen L, Chisholm AD (2011) Axon regeneration mechanisms: insights from *C. elegans*. *Trends Cell Biol* 21:577–584. doi: 10.1016/j.tcb.2011.08.003
57. Chisholm AD (2013) Cytoskeletal Dynamics in *Caenorhabditis elegans* Axon Regeneration. <http://dxdoi.org/101146/annurev-cellbio-101512-122311> 29:271–297. doi: 10.1146/annurev-cellbio-101512-122311
58. Hammarlund M, Nix P, Hauth L, et al (2009) Axon regeneration requires a conserved MAP kinase pathway. *Science* 323:802–

806. doi: 10.1126/science.1165527
59. Nix P, Hisamoto N, Matsumoto K, Bastiani M (2011) Axon regeneration requires coordinate activation of p38 and JNK MAPK pathways. *Proc Natl Acad Sci USA* 108:10738–10743. doi: 10.1073/pnas.1104830108
 60. Yan D, Wu Z, Chisholm AD, Jin Y (2009) The DLK-1 Kinase Promotes mRNA Stability and Local Translation in *C. elegans* Synapses and Axon Regeneration. *Cell* 138:1005–1018. doi: 10.1016/j.cell.2009.06.023
 61. Byrne AB, Walradt T, Gardner KE, et al (2014) Insulin/IGF1 signaling inhibits age-dependent axon regeneration. *Neuron* 81:561–573. doi: 10.1016/j.neuron.2013.11.019
 62. Li C, Hisamoto N, Nix P, et al (2012) The growth factor SVH-1 regulates axon regeneration in *C. elegans* via the JNK MAPK cascade. *Nat Neurosci* 15:551–557. doi: 10.1038/nn.3052
 63. Yan D, Jin Y (2012) Regulation of DLK-1 Kinase Activity by Calcium-Mediated Dissociation from an Inhibitory Isoform. *Neuron* 76:534–548.
 64. Ghosh-Roy A, Goncharov A, Jin Y, Chisholm AD (2012) Kinesin-13 and Tubulin Posttranslational Modifications Regulate Microtubule Growth in Axon Regeneration. *Dev Cell* 23:716–728.
 65. Kirszenblat L, Neumann B, Coakley S (2013) A dominant mutation in *mec-7/β-tubulin* affects axon development and regeneration in *Caenorhabditis elegans* neurons. *Molecular biology of ...*
 66. Zou Y, Chiu H, Zinovyeva A, et al (2013) Developmental Decline in Neuronal Regeneration by the Progressive Change of Two Intrinsic Timers. *Science* 340:372–376. doi: 10.1126/science.1231321
 67. Kaletsky R, Lakhina V, Arey R, et al (2016) The *C. elegans* adult neuronal IIS/FOXO transcriptome reveals adult phenotype regulators. *Nature* 529:92–96. doi: 10.1038/nature16483
 68. Chiu H, Alqadah A, Chuang C-F, Chang C (2011) *C. elegans* as a genetic model to identify novel cellular and molecular mechanisms underlying nervous system regeneration. *Cell Adh Migr* 5:387–394. doi: 10.4161/cam.5.5.17985

69. Park KK, Liu K, Hu Y, et al (2008) Promoting Axon Regeneration in the Adult CNS by Modulation of the PTEN/mTOR Pathway. *Science* 322:963–966. doi: 10.1126/science.1161566
70. Christie KJ, Webber CA, Martinez JA, et al (2010) PTEN inhibition to facilitate intrinsic regenerative outgrowth of adult peripheral axons. *J Neurosci* 30:9306–9315. doi: 10.1523/JNEUROSCI.6271-09.2010
71. Geoffroy CG, Hilton BJ, Tetzlaff W, Zheng B (2016) Evidence for an Age-Dependent Decline in Axon Regeneration in the Adult Mammalian Central Nervous System. *Cell Rep* 15:238–246. doi: 10.1016/j.celrep.2016.03.028
72. Wood WB (1987) *The nematode Caenorhabditis elegans*. Cold Spring Harbour Laboratory
73. Hart A (2006) Behavior. *WormBook*, ed. The *C. elegans* Research Community, *WormBook*, doi/10.1895/wormbook.1.7.1. doi: 10.1895/wormbook
74. Collins JJ, Huang C, Hughes S, Kornfeld K (2008) The measurement and analysis of age-related changes in *Caenorhabditis elegans*. *WormBook* 1–21. doi: 10.1895/wormbook.1.137.1
75. Croll NA, Smith JM, Zuckerman BM (2007) The aging process of the nematode *Caenorhabditis elegans* in bacterial and axenic culture. *Experimental Aging Research* 3:175–189. doi: 10.1080/03610737708257101
76. Bolanowski MA, Russell RL, Jacobson LA (1981) Quantitative measures of aging in the nematode *Caenorhabditis elegans*. I. Population and longitudinal studies of two behavioral parameters. *Mech Ageing Dev* 15:279–295.
77. Duhon SA, Johnson TE (1995) Movement as an Index of Vitality: Comparing Wild Type and the age-1 Mutant of *Caenorhabditis elegans*. *J Gerontol A Biol Sci Med Sci* 50A:B254–B261. doi: 10.1093/gerona/50A.5.B254
78. Wolkow CA (2006) Identifying factors that promote functional aging in *Caenorhabditis elegans*. *Exp Gerontol* 41:1001–1006. doi: 10.1016/j.exger.2006.06.033
79. Johnson TE (1987) Aging can be genetically dissected into component processes using long-lived lines of *Caenorhabditis ele-*

- gans. *Proc Natl Acad Sci USA* 84:3777–3781.
80. Huang C, Xiong C, Kornfeld K (2004) Measurements of age-related changes of physiological processes that predict lifespan of *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 101:8084–8089. doi: 10.1073/pnas.0400848101
 81. Glenn CF, Chow DK, David L, et al (2004) Behavioral deficits during early stages of aging in *Caenorhabditis elegans* result from locomotory deficits possibly linked to muscle frailty. *J Gerontol A Biol Sci Med Sci* 59:1251–1260.
 82. Mulcahy B, Holden-Dye L, O'Connor V (2013) Pharmacological assays reveal age-related changes in synaptic transmission at the *Caenorhabditis elegans* neuromuscular junction that are modified by reduced insulin signalling. *216:492–501*. doi: 10.1242/jeb.068734
 83. García LR (2014) Regulation of sensory motor circuits used in *C. elegans* male intromission behavior. *Semin Cell Dev Biol* 33:42–49. doi: 10.1016/j.semcd.2014.05.006
 84. Guo X, García LR (2014) SIR-2.1 integrates metabolic homeostasis with the reproductive neuromuscular excitability in early aging male *Caenorhabditis elegans*. *Elife* 3:e01730. doi: 10.7554/eLife.01730
 85. Guarente L (2001) SIR2 and aging – the exception that proves the rule. *Trends in Genetics* 17:391–392.
 86. Ben-Yakar A, Chronis N, Lu H (2009) Microfluidics for the analysis of behavior, nerve regeneration, and neural cell biology in *C. elegans*. *Curr Opin Neurobiol* 19:561–567. doi: 10.1016/j.conb.2009.10.010
 87. Stirman JN, Brauner M, Gottschalk A, Lu H (2010) High-throughput study of synaptic transmission at the neuromuscular junction enabled by optogenetics and microfluidics. *J Neurosci Methods* 191:90–93. doi: 10.1016/j.jneumeth.2010.05.019
 88. Chokshi TV, Bazopoulou D, Chronis N (2010) An automated microfluidic platform for calcium imaging of chemosensory neurons in *Caenorhabditis elegans*. *Lab Chip* 10:2758–2763. doi: 10.1039/c004658b
 89. Leinwand SG, Yang CJ, Bazopoulou D, et al (2015) Circuit

- mechanisms encoding odors and driving aging-associated behavioral declines in *Caenorhabditis elegans*. *Elife* 4:e10181. doi: 10.7554/eLife.10181
90. Kato S, Kaplan HS, Schrödel T, et al (2015) Global brain dynamics embed the motor command sequence of *Caenorhabditis elegans*. *Cell* 163:656–669. doi: 10.1016/j.cell.2015.09.034
91. Nguyen JP, Shipley FB, Linder AN, et al (2016) Whole-brain calcium imaging with cellular resolution in freely behaving *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 113:E1074–81. doi: 10.1073/pnas.1507110112
92. Hedden T, Gabrieli JDE (2004) Insights into the ageing mind: a view from cognitive neuroscience. *Nat Rev Neurosci* 5:87–96. doi: 10.1038/nrn1323
93. Yeoman M, Scutt G, Faragher R (2012) Insights into CNS ageing from animal models of senescence. *Nat Rev Neurosci* 13:435–445. doi: 10.1038/nrn3230
94. Ardiel EL, Rankin CH (2010) An elegant mind: learning and memory in *Caenorhabditis elegans*. *Learn Mem* 17:191–201. doi: 10.1101/lm.960510
95. Kimata T, Sasakura H, Ohnishi N, et al (2012) Thermotaxis of *C. elegans* as a model for temperature perception, neural information processing and neural plasticity. *Worm* 1:31–41. doi: 10.4161/worm.19504
96. Nishida Y, Sugi T, Nonomura M, Mori I (2011) Identification of the AFD neuron as the site of action of the CREB protein in *Caenorhabditis elegans* thermotaxis. *EMBO Rep* 12:855–862. doi: 10.1038/embor.2011.120
97. Murakami S, Murakami H (2005) The effects of aging and oxidative stress on learning behavior in *C. elegans*. *Neurobiol Aging* 26:899–905. doi: 10.1016/j.neurobiolaging.2004.08.007
98. Murakami H, Bessinger K, Hellmann J, Murakami S (2005) Aging-dependent and -independent modulation of associative learning behavior by insulin/insulin-like growth factor-1 signal in *Caenorhabditis elegans*. *J Neurosci* 25:10894–10904. doi: 10.1523/JNEUROSCI.3600-04.2005
99. Kodama E, Kuhara A, Mohri-Shiomi A, et al (2006) Insulin-like

- signaling and the neural circuit for integrative behavior in *C. elegans*. *Genes Dev* 20:2955–2960. doi: 10.1101/gad.1479906
100. Wang Y, Hekimi S (2015) Mitochondrial dysfunction and longevity in animals: Untangling the knot. *Science* 350:1204–1207. doi: 10.1126/science.aac4357
 101. Feng J, Bussi ere F, Hekimi S (2001) Mitochondrial electron transport is a key determinant of life span in *Caenorhabditis elegans*. *Dev Cell* 1:633–644.
 102. Adachi H, Fujiwara Y, Ishii N (1998) Effects of Oxygen on Protein Carbonyl and Aging in *Caenorhabditis elegans* Mutants With Long (age-1) and Short (mev-1) Life Spans. *J Gerontol A Biol Sci Med Sci* 53A:B240–B244. doi: 10.1093/gerona/53A.4.B240
 103. Ishii N, Fujii M, Hartman PS, et al (1998) A mutation in succinate dehydrogenase cytochrome b causes oxidative stress and ageing in nematodes. *Nature* 394:694–697. doi: 10.1038/29331
 104. Bargmann CI (2006) Chemosensation in *C. elegans* (October 25, 2006), *WormBook*, ed. The *C. elegans* Research Community, *WormBook*, doi/10.1895/wormbook. 1.123. 1.
 105. Kauffman AL, Ashraf JM, Corces-Zimmerman MR, et al (2010) Insulin signaling and dietary restriction differentially influence the decline of learning and memory with age. *PLoS Biol* 8:e1000372. doi: 10.1371/journal.pbio.1000372
 106. Beck CD, Rankin CH (1993) Effects of aging on habituation in the nematode *Caenorhabditis elegans*. *Behav Processes* 28:145–163. doi: 10.1016/0376-6357(93)90088-9
 107. Timbers TA, Giles AC, Ardiel EL, et al (2013) Intensity discrimination deficits cause habituation changes in middle-aged *Caenorhabditis elegans*. *Neurobiol Aging* 34:621–631. doi: 10.1016/j.neurobiolaging.2012.03.016
 108. Pereira S, van der Kooy D (2012) Two forms of learning following training to a single odorant in *Caenorhabditis elegans* AWC neurons. *J Neurosci* 32:9035–9044. doi: 10.1523/JNEUROSCI.4221-11.2012
 109. Stein GM, Murphy CT (2012) The Intersection of Aging, Longevity Pathways, and Learning and Memory in *C. elegans*. *Front Genet* 3:259. doi: 10.3389/fgene.2012.00259

110. Brightwell J (2004) Hippocampal CREB1 but not CREB2 is decreased in aged rats with spatial memory impairments. *Neurobiology of Learning and Memory* 81:19–26. doi: 10.1016/j.nlm.2003.08.001
111. Mouravlev A, Dunning J, Young D, During MJ (2006) Somatic gene transfer of cAMP response element-binding protein attenuates memory impairment in aging rats. *Proc Natl Acad Sci USA* 103:4705–4710. doi: 10.1073/pnas.0506137103
112. Kandel ER (2001) The Molecular Biology of Memory Storage: A Dialogue Between Genes and Synapses. *Science* 294:1030–1038. doi: 10.1126/science.1067020
113. Salih DAM, Rashid AJ, Colas D, et al (2012) FoxO6 regulates memory consolidation and synaptic function. *Genes Dev* 26:2780–2801. doi: 10.1101/gad.208926.112
114. Singh-Manoux A, Kivimaki M, Glymour MM, et al (2012) Timing of onset of cognitive decline: results from Whitehall II prospective cohort study. *BMJ* 344:d7622–d7622. doi: 10.1136/bmj.d7622
115. Steculorum SM, Solas M, Brüning JC (2014) The paradox of neuronal insulin action and resistance in the development of aging-associated diseases. *Alzheimers Dement* 10:S3–11. doi: 10.1016/j.jalz.2013.12.008
116. Prehn K, Jumpertz von Schwartzberg R, Mai K, et al (2016) Caloric Restriction in Older Adults—Differential Effects of Weight Loss and Reduced Weight on Brain Structure and Function. *Cereb Cortex* bhw008. doi: 10.1093/cercor/bhw008
117. Bischof GN, Park DC (2015) Obesity and Aging: Consequences for Cognition, Brain Structure, and Brain Function. *Psychosomatic Medicine* 77:697–709. doi: 10.1097/PSY.0000000000000212
118. Miller AA, Spencer SJ (2014) Obesity and neuroinflammation: A pathway to cognitive impairment. *Brain, Behavior, and Immunity* 42:10–21. doi: 10.1016/j.bbi.2014.04.001
119. Witte AV, Fobker M, Gellner R, et al (2009) Caloric restriction improves memory in elderly humans. *Proc Natl Acad Sci USA* 106:1255–1260. doi: 10.1073/pnas.0808587106
120. Kyriazis M (2009) Calorie restriction mimetics: Examples and mode of action. *Open Longev Sci*



121. Witte AV, Kerti L, Margulies DS, Flöel A (2014) Effects of resveratrol on memory performance, hippocampal functional connectivity, and glucose metabolism in healthy older adults. *J Neurosci* 34:7862–7870. doi: 10.1523/JNEUROSCI.0385-14.2014
122. Onken B, Driscoll M (2010) Metformin Induces a Dietary Restriction–Like State and the Oxidative Stress Response to Extend *C. elegans* Healthspan via AMPK, LKB1, and SKN-1. *PLoS ONE* 5:e8758. doi: 10.1371/journal.pone.0008758
123. de Lau LML, Giesbergen PCLM, De Rijk MC, et al (2004) Incidence of parkinsonism and Parkinson disease in a general population: the Rotterdam Study. *Neurology* 63:1240–1244. doi: 10.1212/01.WNL.0000140706.52798.BE
124. Niccoli T, Partridge L (2012) Ageing as a Risk Factor for Disease. *Current Biology* 22:R741–R752.
125. Bargmann CI, Hartweg E, Horvitz HR (1993) Odorant-selective genes and neurons mediate olfaction in *C. elegans*. *Cell* 74:515–527. doi: 10.1016/0092-8674(93)80053-H
126. Avery L, Horvitz HR (1989) Pharyngeal pumping continues after laser killing of the pharyngeal nervous system of *C. elegans*. *Neuron* 3:473–485. doi: 10.1016/0896-6273(89)90206-7
127. Dimitriadi M, Hart AC (2010) Neurodegenerative disorders: Insights from the nematode *Caenorhabditis elegans*. *Neurobiol Dis* 40:4–11. doi: 10.1016/j.nbd.2010.05.012
128. Li J, Le W (2013) Modeling neurodegenerative diseases in *Caenorhabditis elegans*. *Exp Neurol* 250:94–103. doi: 10.1016/j.expneurol.2013.09.024
129. Markaki M, Tavernarakis N (2010) Modeling human diseases in *Caenorhabditis elegans*. *Biotechnology Journal* 5:1261–1276. doi: 10.1002/biot.201000183
130. Harrington AJ, Hamamichi S, Caldwell GA, Caldwell KA (2010) *C. elegans* as a model organism to investigate molecular pathways involved with Parkinson's disease. *Dev Dyn* 239:1282–1295. doi: 10.1002/dvdy.22231
131. Caldwell GA, Caldwell KA (2008) Traversing a wormhole to combat Parkinson's disease. *Dis Model Mech* 1:32–36. doi: 10.1242/dmm.000257

132. Hannan SB, Dräger N, Rasse TM, et al (2016) Cellular and molecular modifier pathways in tauopathies: the big picture from screening invertebrate models. *J Neurochem* n/a–n/a. doi: 10.1111/jnc.13532
133. Link CD (2006) *C. elegans* models of age-associated neurodegenerative diseases: Lessons from transgenic worm models of Alzheimer's disease. *Exp Gerontol* 41:1007–1013. doi: 10.1016/j.exger.2006.06.059
134. Lublin AL, Link CD (2013) Alzheimer's disease drug discovery: in vivo screening using *Caenorhabditis elegans* as a model for β -amyloid peptide-induced toxicity. *Drug Discovery Today: Technologies* 10:e115–e119. doi: 10.1016/j.ddtec.2012.02.002
135. Pouladi MA, Morton AJ, Hayden MR (2013) Choosing an animal model for the study of Huntington's disease. *Nat Rev Neurosci* 14:708–721. doi: 10.1038/nrn3570
136. Therrien M, Parker JA (2014) Worming forward: amyotrophic lateral sclerosis toxicity mechanisms and genetic interactions in *Caenorhabditis elegans*. *Front Genet* 5:85. doi: 10.3389/fgene.2014.00085
137. Nikolettou V, Tavernarakis N (2014) Necrotic cell death in *Caenorhabditis elegans*. *Meth Enzymol* 545:127–155. doi: 10.1016/B978-0-12-801430-1.00006-8
138. Mano I, Driscoll M (2009) *Caenorhabditis elegans* glutamate transporter deletion induces AMPA-receptor/adenylyl cyclase 9-dependent excitotoxicity. *J Neurochem* 108:1373–1384. doi: 10.1111/j.1471-4159.2008.05804.x
139. Nagarajan A, Ning Y, Reisner K, et al (2014) Progressive degeneration of dopaminergic neurons through TRP channel-induced cell death. *J Neurosci* 34:5738–5746. doi: 10.1523/JNEUROSCI.4540-13.2014
140. Hamamichi S, Rivas RN, Knight AL, et al (2008) Hypothesis-based RNAi screening identifies neuroprotective genes in a Parkinson's disease model. *Proc Natl Acad Sci USA* 105:728–733. doi: 10.1073/pnas.0711018105
141. Gitler AD, Chesi A, Geddie ML, Strathearn KE (2009) α -Synuclein is part of a diverse and highly conserved interaction network that includes PARK9 and manganese toxicity. *Nature*

142. Qiao L, Hamamichi S, Caldwell KA, et al (2008) Lysosomal enzyme cathepsin D protects against alpha-synuclein aggregation and toxicity. *Molecular Brain* 2008 1:1 1:1. doi: 10.1186/1756-6606-1-17
143. Knight AL, Yan X, Hamamichi S, et al (2014) The glycolytic enzyme, GPI, is a functionally conserved modifier of dopaminergic neurodegeneration in Parkinson's models. *Cell Metab* 20:145–157. doi: 10.1016/j.cmet.2014.04.017
144. Kraemer BC, Burgess JK, Chen JH, et al (2006) Molecular pathways that influence human tau-induced pathology in *Caenorhabditis elegans*. *Hum Mol Genet* 15:1483–1496. doi: 10.1093/hmg/ddl067
145. Amaducci L, Tesco G (1994) Aging as a major risk for degenerative diseases of the central nervous system. *Curr Opin Neurol* 7:283–286.
146. Volovik Y, Marques FC, Cohen E (2014) The nematode *Caenorhabditis elegans*: a versatile model for the study of proteotoxicity and aging. *Methods* 68:458–464. doi: 10.1016/j.ymeth.2014.04.014
147. Morley JF, Brignull HR, Weyers JJ, Morimoto RI (2002) The threshold for polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and influenced by aging in *Caenorhabditis elegans*. *Proc Natl Acad Sci USA* 99:10417–10422. doi: 10.1073/pnas.152161099
148. Hsu A-L, Murphy CT, Kenyon C (2003) Regulation of aging and age-related disease by DAF-16 and heat-shock factor. *Science* 300:1142–1145. doi: 10.1126/science.1083701
149. Cohen E, Bieschke J, Perciavalle RM, et al (2006) Opposing activities protect against age-onset proteotoxicity. *Science* 313:1604–1610. doi: 10.1126/science.1124646
150. Florez-McClure ML, Hohsfield LA, Fonte G, et al (2007) Decreased insulin-receptor signaling promotes the autophagic degradation of beta-amyloid peptide in *C. elegans*. *Autophagy* 3:569–580.
151. Zhang T, Mullane PC, Periz G, Wang J (2011) TDP-43 neurotoxicity and protein aggregation modulated by heat shock factor and insulin/IGF-1 signaling. *Hum Mol Genet* 20:1952–1965. doi:

- 10.1093/hmg/ddr076
152. David DC, Ollikainen N, Trinidad JC, et al (2010) Widespread protein aggregation as an inherent part of aging in *C. elegans*. *PLoS Biol* 8:e1000450. doi: 10.1371/journal.pbio.1000450
 153. Steinkraus KA, Smith ED, Davis C, et al (2008) Dietary restriction suppresses proteotoxicity and enhances longevity by an hsf-1-dependent mechanism in *Caenorhabditis elegans*. *Aging Cell* 7:394–404. doi: 10.1111/j.1474-9726.2008.00385.x
 154. Jadiya P, Chatterjee M, Sammi SR, et al (2011) Sir-2.1 modulates 'calorie-restriction-mediated' prevention of neurodegeneration in *Caenorhabditis elegans*: implications for Parkinson's disease. *Biochem Biophys Res Commun* 413:306–310. doi: 10.1016/j.bbrc.2011.08.092
 155. Shemesh N, Shai N, Ben-Zvi A (2013) Germline stem cell arrest inhibits the collapse of somatic proteostasis early in *Caenorhabditis elegans* adulthood. *Aging Cell* 12:814–822. doi: 10.1111/accel.12110
 156. Cohen E, Paulsson JF, Blinder P, et al (2009) Reduced IGF-1 signaling delays age-associated proteotoxicity in mice. *Cell* 139:1157–1169. doi: 10.1016/j.cell.2009.11.014
 157. Gontier G, George C, Chaker Z, et al (2015) Blocking IGF Signaling in Adult Neurons Alleviates Alzheimer's Disease Pathology through Amyloid- β Clearance. *J Neurosci* 35:11500–11513. doi: 10.1523/JNEUROSCI.0343-15.2015
 158. De Magalhaes Filho CD, Kappeler L, Dupont J, et al (2016) Deleting IGF-1 receptor from forebrain neurons confers neuroprotection during stroke and upregulates endocrine somatotropin. *J Cereb Blood Flow Metab* 0271678X15626718. doi: 10.1177/0271678X15626718
 159. Wang J, Ho L, Qin W, et al (2005) Caloric restriction attenuates beta-amyloid neuropathology in a mouse model of Alzheimer's disease. *FASEB J* 19:659–661. doi: 10.1096/fj.04-3182fje
 160. Maswood N, Young J, Tilmont E, et al (2004) Caloric restriction increases neurotrophic factor levels and attenuates neurochemical and behavioral deficits in a primate model of Parkinson's disease. *Proc Natl Acad Sci USA* 101:18171–18176. doi: 10.1073/pnas.0405831102

161. Cohen E, Du D, Joyce D, et al (2010) Temporal requirements of insulin/IGF-1 signaling for proteotoxicity protection. *Aging Cell* 9:126–134. doi: 10.1111/j.1474-9726.2009.00541.x



| Genotype Phenotype | Wild type | Insulin signalling | | <i>hsf-1</i> | Dietary restriction | Mitochondria | |
|--|---|--|--|---|--|---|--|
| | | <i>daf-2</i> | <i>daf-16</i> | | <i>eat-2</i> | <i>clk-1</i> | |
| Neuronal morphology (Touch neurons) Abnormal branching, shape, wavy axons defasciculation | Appearance of neuron-specific changes with age [6-8] | In some cases decreased appearance of morphological changes with age [6-8] | Increased appearance of morphological changes [7] Suppresses <i>daf-2</i> effects [8] | Increased changes [6-8] | In some cases similar to WT [8] | Decreased morphological changes [8] | |
| | | In other cases increased [7,9] | In other cases, decreased [7,9] | | In other cases decreased [9] | | |
| Synapses Number of synaptic vesicles and puncta Amplitude of post-synaptic currents PSCs | Vesicle and puncta decline (d15, 18) [6, 11] PSC decline (starting d7) [13] | Synaptic puncta maintained (d18, d30) [11] PSC maintenance (up to d27) [13] | | | | | |
| Mitochondria (in ALM) Mitochondrial load | Increases (up to d4) Maintained (d4-8), Decreases (after d8) [36] | Lower load than WT, but steady (d1 to 25) [36] | Similar to WT [36] | | Lower than WT (d4-8), and steady (d4 to d11) [36] | | |
| Mitochondrial transport | Decreases after d1 [36] | Steady (d1 to 25) [36] | Similar to WT [36] | | Steady (d1 to 11) [36] | | |
| Resistance to oxidative stress | Increases (up to d4) Decreases after d4 [36] | Higher resistance, lower rate of decline until d22 [36] | | | | | |
| Axon regeneration GABA motor neurons | Declines from d1 Abolished by d5 [5] | Delayed decline (no decline on d5, decline by d10) [61] | Suppresses increased regeneration of <i>daf-2</i> [61] | | Similar to WT [61] | | |
| Learning and Memory Thermotaxis learning | Declines (d6) Absent (d11) [98] | Enhanced learning in young Delayed decline in old [98] | Suppresses <i>daf-2</i> delayed decline [97, 98] | | Increased learning in young. Delayed decline with age [98] | Enhanced learning in young. Delayed decline in old [97, 98] | |
| | | Longer in young animals (40 vs 24 hr in the WT) [11] Not extended in aged animals [105] | | | Defective in LTAM [105] | Impaired in young adults | |
| | | Improved in older animals [105] | | | | | |
| STAM (positive olfactory) | Massed learning: decline begins d3, abolished by d6. Spaced learning lasts until d7 [105] | 3x longer STAM in young adults Maintained in older worms (no loss in d5) [105] | Defective in STAM, suppresses <i>daf-2</i> STAM extension [11, 105] | | Similar to WT in young adults [105] Improved in older animals (after spaced learning) [105] | | |
| Neurodegeneration, proteotoxicity Aggregation of PolyQ, A β , SOD-1, α -synuclein | Increased aggregation / proteotoxicity with age | Reduced aggregation and proteotoxicity [143, 147-151] | Suppresses <i>daf-2</i> protective effect [143, 147-151] | Required for <i>daf-2</i> and dietary restriction protective effects [143, 148-151] | Protects from proteotoxicity [153] | | |